Application of Plasma Phenomena



Po-Yu Chang

Institute of Space and Plasma Sciences, National Cheng Kung University

Lecture 13

2025 fall semester

Thursday 9:00-12:00

Materials:

https://capst.ncku.edu.tw/PGS/index.php/teaching/

Online courses:

https://reurl.cc/ZIG6k6

Grading




- Quizzes 50 % (2-min Q&A at the beginning of each class)
- Presentations 50 % (10-min presentation on any plasma applications or phenomena)
- Prerecorded classes:
 - 1/8: International workshop on multiscale laboratory astrophysics with the aid of data science

Final presentation on 12/31 at 10:30 in room 48111B



Please select all the times you are available so we can find a time slot that works for everyone. 6 則回應



Note!



Final presentation on 12/31 at 10:30 in room 48111B!

Time: 12/31 10:30

Time for each person: 10 mins

- Topics: anything related to plasma. You can talk about a plasma phenomenon or a plasma application. In your presentation, you need to describe how plasma is involved. What rule does the plasma play in the phenomenon or application.
- Your scores (34 points in total) will depend on:

(2 points) Presentation time.

< 8 min: 0

8~9 min: 1

9~11 min: 2

11~12 min: 1


> 12 min: 0

- (2 points) Plasma related.
- (10 points) How clear the physics is explained?
- (10 points) The presentation skill.
- (10 points) Your slides.

EUV light sources


A semiconductor device is fabricated by many repetitive production process





Ultraviolet lithography (EUVL) is one of the key technologies in semiconductor manufacturing nowadays

 The process technology of Taiwan Semiconductor Manufacturing Company Limited (TSMC):



- Optical diffraction needs to be taken into account.
- Shorter wavelength is preferred.
 - Light source with a center wavelength of 13.5 nm is used.

EUV lithography becomes important for semiconductor industry




0.15 billion USD for each EUV light source.

https://www.youtube.com/watch?v=NHSR6AHNiDs

EUV light can only be reflected using multilayer mirrors






https://henke.lbl.gov/optical_constants/ Mo/Si multilayer coating technology for EUVL, coating uniformity and time stability; E. Louis et al.; SPIE 4146-06, Soft X-ray and EUV Imaging Systems, San Diego, 2000.

13.5-nm EUV light is picked for EUV lithography






- $\lambda = 13.5 \text{ nm} \pm 1\%$ is required.
- At T=35-40 eV (~450,000 K), in-band emission occurs.
- Xenon:
 - $4p^64d^8 \rightarrow 4p^64d^75p$ from single ion stage Xe^{10+}
 - UTA @ 11 nm

- Tin:
 - $4p^64d^N \rightarrow 4p^54d^{N+1} + 4p^64d^{N-1}4f$ (1 \leq N \leq 6) in ions ranging from Sn⁸⁺ to Sn¹²⁺
 - UTA @ 13.5 nm
 - UTA: unresolved transition array


EUV light is generated from laser-produced plasma (LPP)






Two laser pulses are used to heat the plasma









Hydrogen buffer gas with a pressure of ~100 Pa is used to protect the collector mirror





Laser-produced plasma (LPP) is used in the EUV lithography





High harmonic generation from high-power laser

Distance

For $I < 10^{18} \text{ w/cm}^2$ Xe Source gas cell IR field a b C (a) atomic core atomic core (b) 3 E_{rad} half optical period = π/ω Energy $\Lambda\Lambda\Lambda\Lambda\Lambda\Lambda\Lambda$ (c) $S(\omega)$ twice laser frequency = $2\omega_{L}$ $q_{\text{max}} \omega_{\text{L}}$


M. Krüger, etc., Appl. Sci. 9, 378 (2019)

Nonlinear Optics 3rd edition, by Robert W. Boyd

P. B. Corkum and F. Krausz, Nature Phys., 3, 381 (2007)


EUV light can be generated using discharged-produced plasma








$$TV^{\gamma-1} = \text{const} \quad T_{\rm f} = T_{\rm o} \left(\frac{r_{\rm o}}{r_{\rm f}}\right)^{4/3}$$




Light source and display systems



Plasma display panel (PDP)



Liquid crystal display (LCD)



Outlines




- Cathode Ray Tube
- Color space (CIE 1931 color spaces)
- History of plasma display panel (PDP)
- Design of PDP
- Liquid crystal display (LCD)
- LCD vs PDP




Cathode Ray Tube uses electron beams to light the fluorescent screen







The image is shown by scanning through the whole screen with the single electron beam










Color image is formed by using three electron beams scanning through three different color channels






Color space


Color can be created using three primary colors






Human retina has three kinds of "cones" that have different spectral response






Spectral response of retina "cones" are tested using light sources with single wavelength





The CIE 1931 color space chromaticity diagram is the standard color space





0.1

0.2

0.3

0.4


0.5

0.6

0.7

History of PDP

Plasma display panel was invented at the University of Illinois in 1967



Prof. H. Gene Slottow

Prof. Donald L. Bitzer

PDP was invented due to a need for Programmed Logic for Automatic Teaching Operations (PLATO) in 1960s











https://topwallpapers.pw/computer/keyboards-computers-history-teletype-typewriters-desktop-hd-wallpaper-1035981/https://en.wikipedia.org/wiki/Punched_tape https://en.wikipedia.org/wiki/PLATO (computer system)

The positive column in a glow discharge is used to excite phosphors in color PDP








- Majority of monochrome PDPs use the negative glow as the light source
- The positive column is used to excite phosphors in fluorescent lamps and in color PDPs

Early plasma panel (PD) attached to the glass vacuum system used for the first plasma displays at UI






 It had the same alternating sustain voltage, neon, gas, and dielectric glass insulated electrodes that are used for plasma TVs today.

Early plasma panel (PD) attached to the glass vacuum system used for the first plasma displays at UI






 It had the same alternating sustain voltage, neon, gas, and dielectric glass insulated electrodes that are used for plasma TVs today.


Early 4x4 pixel panel has achieved matrix addressability for the first time






Early 4x4 pixel panel has achieved matrix addressability for the first time






A 16x16 pixel PD, developed in 1967, needed to be addressed manually






First color PD was three cell prototype with red and green color phosphors excited by a xenon gas discharge



Open-cell structure developed in 1968








It could be baked under vacuum at 350 °C to drive out contaminants.

More progress




1968, University of Illinois 16x16 pixels

1971, Owens-Illinois 512x512 pixels






Color PDPs had short display lifetime due to the degradation of color phosphors caused by ion sputtering




Design of PDP

A lower breakdown voltages can be obtained with very small amounts of added gas




AT&T three-electrode patent







Reflective phosphor geometry is used in most of today's plasma TVs











Spectrum of the different phosphors






The foundation of AC discharge








The plasma can be sustained using ac discharged






Wall discharge reduced the required discharge voltage


Wall discharge reduced the required discharge voltage





ON/OFF State Selection








Sustain discharge




ON Cell













OFF Cell










Address and sustain electrodes are connected to different drivers







PDP pixel can only be either ON or OFF



Cathode Ray Tube :



Plasma Display Panel :



PDP luminance is controlled by using number of light pulses



CRT : Control the Luminance using Electron Beam Intensity




PDP : Control the Luminance using Number of Light Pulses




A single field is divided into 8 subfield





Composition of each subfield






Spec: VGA (640*480) 8 Subfield

0.03msec Address Pulse

100KHz Sustain Freq.

8 subfield in one TV-Field (ADS)







PDP uses line-by-line scanning



Cathode Ray Tube : Cell-by-Cell Scanning




PDP : Line-by-Line Scanning



Video signal processing



Analog Video Signal ⇒ Digital Pulse Signal



Addressing period





Displaying period









Liquid crystal are a special state of matter between liquid and crystal






Linear polarization of a light can be rotated by miss aligned liquid crystal






Structure of Liquid crystal display (LCD)





Optimistic projection of PDP market






Reality



TV Shipment Growth by Technology




Too many reasons that PDP died!



- Bright showroom conditions put plasmas at a distinct disadvantage versus LED-lit LCDs
- Aesthetics may have played a role in hastening plasma's demise
- UHD/4K caught on quickly
- Screen-size limitations also played a part in plasmas plight
- You can't bend a plasma
- Plasmas were harder to deal with than LCDs
- While OLED is still in the early stages of development, there's no question it offers greater potential than plasma
- Energy efficiency may have played a part in putting plasma out to pasture
- Plasma was the original flat-panel technology, People just thought of it as old technology.
- Projectors improved in quality and prices dropped

Let's stand up and do exercise!!






The hydrogen bomb






The "iron group" of isotopes are the most tightly bound





Chain reaction can happen in U²³⁵ fission reaction






- ~ 200 million electron volts (MeV)/fission, ~million times more than chemical reactions
- Energy for bombs, or for civilian power can generate huge amounts of energy (and toxicity) in a small space with a modest amount of material
- Source of safety, security issues for nuclear power


The neutrons are leaking out and stopping the chain reaction in a sub-critical mass






Solution 1: add more material






Solution2: reflect the neutron back in






Solution 3: increase the density





How to get the material together before it blows apart?






- There are always neutrons around
- Once chain reaction starts, material will heat up, expand, stop reaction
- How to get enough material together fast enough?

Gun-type bomb




- Simple, reliable can be built without testing
- Highly inefficient require lots of nuclear material (50-60 kg of 90% enriched HEU)
- Can only get high yield with HEU, not plutonium
- Hiroshima bomb: cannon that fired HEU projectile into HEU target



Implosion design



 A schematic diagram of an implosion bomb



 Small-scale slow-motion cross-section of a shaped charge implosion design




The 1st nuclear bomb: Trinity (Bradbury Science Museum)

Model of the Trinity Gadget



 Project Y Atomic Bomb Detonator System



https://www.flickr.com/photos/rocbolt/with/8061684482

Project Y Atomic Bomb Detonator System





 Project Y Atomic Bomb Detonator System Spark Gap Switch



The 1st nuclear bomb: Trinity









https://www.theatlantic.com/photo/2015/07/70-years-since-trinity-when-we-tested-nuclear-bombs/398735/https://saddlebagnotes.com/arts-and-leisure/tucson-seismographs-detected-first-nuclear-test-at-trinity-n-m/article_b01c5b20-f6fb-11eb-a221-6327df2feaeb.html

Trinity explosion on July 16, 1945









https://www.theatlantic.com/photo/2015/07/70-years-since-trinity-when-we-tested-nuclear-bombs/398735/https://en.wikipedia.org/wiki/Trinity_%28nuclear_test%29

Hiroshima Bomb – "Little Boy"





Gun Type – Easiest to design and build (Hiroshima bomb was never tested)

About 13 kiloton explosive yield

Atomic bomb is very destructive



Hiroshima: August 6, 1945




Nagasaki: August 9, 1945



The fusion process





$$^{2}H+^{3}H \Rightarrow ^{4}He+n+Q \equiv 17.6 \text{ MeV}$$

Energy release Q=17.6 MeV

In comparison

$${}^{2}H+{}^{2}H \Rightarrow {}^{1}H+{}^{3}H +Q \equiv 4.0 \text{ MeV}$$
 ${}^{2}H+{}^{2}H \Rightarrow {}^{3}He+n +Q \equiv 3.2 \text{ MeV}$
 ${}^{3}H+{}^{3}H \Rightarrow {}^{4}He+2n+Q \equiv 11.3 \text{ MeV}$
 ${}^{235}U+n \Rightarrow X_{A}+X_{B}+3n +Q \approx 200 \text{ MeV}$

Deuterium-Tritium Fusion Reaction

Fusionable Material, deuterium ²H (D) and tritium ³H (t):

Deuterium: natural occurrence (heavy water) (0.015%).

Tritium: natural occurrence in atmosphere through cosmic ray bombardment; radioactive with $T_{1/2}$ =12.3 y.

"Advantages" of hydrogen bomb



Fusion of
$${}^{2}\text{H+}{}^{3}\text{H}$$
: $\frac{Q}{A} = \frac{17.6 \ MeV}{(3+2) \ amu} = 3.5 \frac{MeV}{amu}$

Fission of ²³⁵U:
$$\frac{Q}{A} = \frac{200 \ MeV}{236 \ amu} = 0.85 \frac{MeV}{amu}$$

Fusion is 4 times more powerful than fission and generates 24 times more neutrons!


$$^{2}H + ^{3}H : \frac{n}{A} = \frac{1}{5} = 0.2$$

Neutron production:

$$^{235}U + n$$
: $\frac{n}{A} = \frac{2}{236} = 0.0085$

Hydrogen bomb uses a fission bomb to initiate the fusion reaction






Fuel

Primary Fission Device

Core: ²³⁹Pu, ²³⁵U, plus ²H+³H booster

Shell: 238U tamper

High explosive lenses




Secondary Fusion Device


Radiation channel
²³⁹Pu sparkplug
⁶Li, ²H, ³H fusion cell
²³⁸U tamper

Event sequence






1. Warhead before firing; primary (fission bomb) at top, secondary (fusion fuel) at bottom, all suspended and beginning a fission in polystyrene foam.



2. HE fires in primary, compressing plutonium core into supercriticality reaction.



 Fissioning primary emits X-rays which reflect along the inside of the casing, irradiating the polystyrene foam.



Polystyrene foam becomes plasma, compressing secondary, and plutonium sparkplug begins to fission.




Compressed and heated, lithium-6 deuteride fuel begins fusion reaction, neutron flux causes tamper to fission. A fireball is starting to form...

Additional pressure from recoil of exploding shell (ablation)!


You don't want to build a hydrogen bomb!






To Fuse, or Not to Fuse...







Outline




- Introduction to nuclear fusion
- Magnetic confinement fusion (MCF)
 - Tokamak
 - Stellarator
- Inertial confinement fusion (ICF)
 - Indirection drive ICF
 - Direct drive ICF
- Innovation idea MCF + ICF
- Plasma in space
- Pulsed-power system at NCKU

World energy consumption is dominated by the use of dwindling fossil fuels



Fossil fuel	Estimated reserve	(2005 consumption rate) Years remaining
Oil	1,277,702 million barrels	32 years
Natural gas	~6,500,000 billion cubic ft	72 years
Coal	1,081,279 million tons	252 years



E15657


While predictions about the exact number of remaining years vary, fossil fuels will run out.

Energy Information Administration (EIA) 2006 Annual Report, U.S. Department of Energy, Washington, D.C.

^{*}from Laboratory for Laser Energetics, University of Rochester, Rochester, NY

Nuclear energy has the highest energy density






Nuclear energy has the highest energy density




HOW MUCH LAND DOES IT TAKE TO POWER A CITY OF 1 MILLION*?



The "iron group" of isotopes are the most tightly bound





$$Q = \left(\sum_{i} m_{i} - \sum_{f} m_{f}\right) c^{2}$$

$$\Delta m = z m_{\rm p} + (A - z) m_{\rm n} - m$$


• Binding energy: $B = \Delta mc^2$

Output energy: $Q = \sum_{f} B_{f} - \sum_{i} B_{i}$


Fusion in the sun provides the energy



Proton-proton chain in sun or smaller




Particles are confined by the gravity.




In heavy sun, the fusion reaction is the CNO cycle






The cross section is a measure of the probability that a specific process will take place in a collision of two particles







Cross section measures the probability per pair of particles for the occurrence of the reaction



$$x_1 + x_2 \rightarrow x_3 + x_4$$

The hard sphere cross section: $\sigma \approx \pi R^2$ where R $\approx 5 \times 10^{-15}$ m is the nuclear radius, i.e., $\sigma = 0.8 \times 10^{-28}$ m² ≈ 1 barm. (barm $\equiv 10^{-28}$ m²)



Classical cross section:


$$D \biguplus v_{D} \bigvee v_{T}$$

$$\frac{m_{\rm D}}{2}v_{\rm D}^2 + \frac{m_{\rm T}}{2}v_{\rm T}^2 \ge \frac{e^2}{4\pi\epsilon_{\rm o}R}$$

• Let
$$v = \left| \overrightarrow{v}_D - \overrightarrow{v}_T \right|$$

$$v_{\mathrm{D}} = \frac{m_{\mathrm{T}}}{m_{\mathrm{D}} + m_{\mathrm{T}}} v \qquad v_{\mathrm{T}} = \frac{m_{\mathrm{D}}}{m_{\mathrm{D}} + m_{\mathrm{T}}} v$$


$$\frac{m_{\rm r}}{2}v^2 \ge \frac{e^2}{4\pi\epsilon_0 R} \qquad m_{\rm r} = \frac{m_{\rm D}m_{\rm T}}{m_{\rm D} + m_{\rm T}}$$



 Classical kinetic energy required for fusion is

$$K_{\rm c.m.} > 288 \, {\rm keV} \, |||$$

Cross section of fusion reaction is much larger than the classical approach



 $D + He^3 \rightarrow He^4 + p$


 $p + B^{11} \rightarrow 3He^4$

$$\begin{array}{c} D+D \rightarrow T+p \\ \rightarrow He^3+n \end{array}$$

$$D + T \rightarrow He^4 + n$$

• Classical kinetic energy required for fusion is
$$K_{\text{c.m.}} > 288 \text{ keV } \text{!!!}$$

- DT cross section has a peak of ~5 barns at 120 keV.
- $\sigma_{DT} \approx 100\sigma_{DD}$ @ 20 keV.



https://i.stack.imgur.com/wXQD5.jpg Santarius, J. F., "Fusion Space Propulsion – A Shorter Time Frame Than You Think", JANNAF, Monterey, 5-8 December 2005.

The cross section of proton-proton chain is much smaller than D T fusion




Reaction	σ _{10 keV} (barn)	σ _{100 keV} (barn)	σ _{max} (barn)	ε _{max} (keV)
D+T→α+n	2.72x10 ⁻²	3.43	5.0	64
D+D→T+p	2.81x10 ⁻⁴	3.3x10 ⁻²	0.06	1250
D+D→³He+n	2.78x10 ⁻⁴	3.7x10 ⁻²	0.11	1750
T+T→α+2n	7.90x10 ⁻⁴	3.4x10 ⁻²	0.16	1000
$D+^3He\rightarrow \alpha+p$	2.2x10 ⁻⁷	0.1	0.9	250
p+ ⁶ Li→α+ ³ He	6x10 ⁻¹⁰	7x10 ⁻³	0.22	1500
$p+^{11}B\rightarrow 3\alpha$	(4.6x10 ⁻¹⁷)	3x10 ⁻⁴	1.2	550
p+p→D+e++v	(3.6x10 ⁻²⁶)	(4.4x10 ⁻²⁵)		
$p+^{12}C\rightarrow^{13}N+\gamma$	(1.9x10 ⁻²⁶)	2.0x10 ⁻¹⁰	1.0x10.4	400
¹² C+ ¹² C (all branches)		(5.0x10 ⁻¹⁰³)		

• "()" are theoretical values while others are measured values.

Enormous fusion fuel can be produced from sea water






What could you do with 1 kg DT?



- 1 kg DT -> 340 Tera joules
 - You can drive your car for ~40,000 km (back and forth between Keelung and Kaoshiung for 50 times).
 - You can keep your furnace running for 8 years.
 - You can blow things up! 1 TJ = 250 tons of TNT.

Nuclear fusion and fission release energy through energetic neutrons



Nuclear fusion provides more energy per atomic mass unit (amu) than nuclear fission




Fusion of
$${}^{2}\text{H+}{}^{3}\text{H}$$
: $\frac{Q}{A} = \frac{17.6 \ MeV}{(3+2) \ amu} = 3.5 \frac{MeV}{amu}$

Fission of ²³⁵U:
$$\frac{Q}{A} = \frac{200 MeV}{236 amu} = 0.85 \frac{MeV}{amu}$$

	Half-life (years)
U235	7.04x10 ⁸
U238	4.47x10 ⁹
•••	
Tritium	12.3


$$n + {}^{6}Li \rightarrow \alpha + T + 4.86 \text{ MeV}$$

 $n + {}^{7}Li \rightarrow \alpha + T + n - 2.87 \text{ MeV}$



Nuclear fusion power plant proposed by Commonwealth Fusion Systems (CFS)






Fusion is much harder than fission



- Fission: $n + {}^{235}_{92}U o {}^{236}_{92}U o {}^{144}_{56}Ba + {}^{89}_{36}Kr + 3n + 177 \text{ MeV}$
- Fusion: $D + T \rightarrow {}_{2}^{4}He(3.5 \text{ MeV}) + n(14.1 \text{ MeV})$




 Beam particles lose their energy before making a fusion reaction, i.e., they only thermalize the fuel. Therefore, beam fusion does NOT work!

Fast neutrons are slowed down due to the collisions

- A moderator is used to slow down fast neutrons but not to absorb neutrons.
- For $m_M \sim m_N$, the energy decrement is higher. Therefore, H slows down neutron most efficiently.

 $m_N m_M$



The best option is the D in the heavy water (D₂O).


	Energy decrement	Neutron scattering cross section (σ_s) (Barns)	Neutron absorption cross section (σ_a) (Barns)
Н	1	49 (H ₂ O)	0.66 (H ₂ O)
D	0.7261	10.6 (D ₂ O)	0.0013 (D ₂ O)
С	0.1589	4.7 (Graphite)	0.0035 (Graphite)

https://en.wikipedia.org/wiki/Neutron_moderator#cite_note-Weston-4 https://energyeducation.ca/encyclopedia/Neutron_moderator#cite_note-3


A temperature describes the randomness of particles

















A "hot plasma" at 100M °C is needed




 Probability for fusion reactions to occur is low at low temperatures due to the coulomb repulsion force.

$$D + \longrightarrow T$$

 If the ions are sufficiently hot, i.e., large random velocity, they can collide by overcoming coulomb repulsion








*R. Betti, HEDSA HEDP Summer School, 2015


https://ydweb.yuda.tyc.edu.tw/Page/YDW/School/Home/Post?id=13286&BoardName=%E6%A1%83%E8%82%B2%E6%B4%BB%E5%8B%95%E7%85%A7%E7%89%87&Board_PartialViewName=&Post_PartialViewName=&Page=60&PageSize=3https://www.freeimages.com/tw/premium/school-playground-with-children-1496555


Fusion doesn't come easy



Probability








$$D + T \rightarrow He^4 + n$$

D + He
3
 \rightarrow He 4 + p

$$p + B^{11} \rightarrow 3He^4$$




Cross section

https://i.stack.imgur.com/wXQD5.jpg Santarius, J. F., "Fusion Space Propulsion - A Shorter Time Frame Than You Think", JANNAF, Monterey, 5-8 December 2005.

Nuclear power plant






Comparison between nuclear fission and nuclear fusion




	Nuclear Fission	Nuclear Fusion
Chain reaction	Yes	No
Melt down	Possible	Impossible
Nuclear waste	High radiative	Low radiative / None

Nuclear fission power plant




Nuclear fussion power plant



It takes a lot of energy or power to keep the plasma at 100M °C



Let the plasma do it itself!



The α-particles heat the plasma.

Under what conditions the plasma keeps itself hot?



Steady state 0-D power balance:

$$S_{\alpha}+S_{h}=S_{B}+S_{k}$$

 S_{α} : α particle heating

S_h: external heating

S_B: Bremsstrahlung radiation

S_k: heat conduction lost


Ignition condition: PT > 10 atm-s = 10 Gbar - ns

- P: pressure, or called energy density
- т is confinement time

The plasma is too hot to be contained




 Solution 1: Magnetic confinement fusion (MCF), use a magnetic field to contain it. P~atm, τ~sec, T~10 keV (10⁸ °C)



Don't confine it!



 Solution 2: Inertial confinement fusion (ICF). Or you can say it is confined by its own inertia: P~Gigabar, τ~nsec, T~10 keV (10⁸ °C)




Inertial confinement fusion: an introduction, Laboratory for Laser Energetics, University of Rochester

To control? Or not to control?




Magnetic confinement fusion (MCF)



Plasma is confined by toroidal magnetic field.

Inertial confinement fusion (ICF)




A DT ice capsule filled with DT gas is imploded by laser.

Laboratory for Laser Energetics, University of Rochester is a pioneer in laser fusion


"Ignition" (target yield larger than one) was achieved for the first time in NIF on 2022/12/5

- Magnetic confinement fusion (MCF)
- Inertial confinement fusion (ICF)



Record-breaking 59
megajoules of sustained
fusion energy in Joint
European Torus (JET) facility
in Oxford demonstrates
powerplant potential and
strengthens case for ITER.





NIF's ignition achievement in perspective

https://ccfe.ukaea.uk/resources/#gallery https://physicstoday.scitation.org/do/10.1063/PT.6.2.20221213a/full/ https://lasers.llnl.gov/science/achieving-fusion-ignition


Outline




- Introduction to nuclear fusion
- Magnetic confinement fusion (MCF)
 - Tokamak
 - Stellarator
- Inertial confinement fusion (ICF)
 - Indirection drive ICF
 - Direct drive ICF
- Innovation idea MCF + ICF
- Plasma in space
- Pulsed-power system at NCKU

Charged particles gyro around the magnetic fields








Charged particles can be partially confined by a magnetic mirror machine




• Charged particles with small $v_{||}$ eventually stop and are reflected while those with large $v_{||}$ escape.



- Large v_{II} may occur from collisions between particles.
- Those confined charged particle are eventually lost due to collisions.

"loffe bars" are added to stabilize the Rayleigh-Taylor instabilities at the center of the mirror machine






A "baseball coil" is obtained if one links the coils and the bars into a single conductor



Baseball coil





MFTF-B mirror machine



Plasma can be confined in a doughnut-shaped chamber with toroidal magnetic field

Tokamak - "toroidal chamber with magnetic coils" (тороидальная камера с магнитными катушками)






https://en.wikipedia.org/wiki/Tokamak#cite_ref-4


Charged particles drift across field lines





A poloidal magnetic field is required to reduce the drift across field lines






A poloidal magnetic field is required to reduce the drift across field lines



The outward force can be compensated by externally applied vertical fields




Externally applied vertical field



Coils in a tokamak






- Toroidal field coils (in poloidal direction) generate toroidal field for confinement.
- Poloidal field coils generate vertical field for plasma positioning and shaping.
- Central solenoid for breakdown and generating plasma current (in toroidal direction) and thus generating poloidal field for confinement.


A divertor is needed to remove impurities and the power that escapes from the plasma





D-shaped tokamak with diverter is more preferred nowadays







Make the plasma closer to the major axis

Spherical tokamak is formed when the aspect ratio of a tokamak is reduced to the order of unity


NSTX @ Princeton



 MegaAmpere Spherical Tokamak (MAST) @ Culham center for fusion energy, UK




ITER ("The Way" in Latin) is one of the most ambitious energy projects in the world today





ITER ("The Way" in Latin) is one of the most ambitious energy projects in the world today

Vacuum vessel




Magnets







Divertor



Cryostat




Blanket



ITER



- T=150M °C
- P=500 MW



ITER – Magnets




• E_B=51 GJ

• T_B=4 K

 Length of Nb₃Sn superconducting strand: 10⁵ km

• B_{T,max}=11.8 T

• B_{P,max}=6 T




ITER - Vacuum vessel



• W = 8000 tons

• $V = 840 \text{ m}^3$

• R = 6 m




ITER - Blanket



440 modules

Thermal load:736 MW




ITER - Divertor



54 cassettes

Thermal load:
 20 MW/m²

Each cassette:10 tons




ITER – Crystat



 $P = 10^{-6}$ atm

• W = 3800 tons

• $V = 16000 \text{ m}^3$




Supporting systems



- Tritium breeding
- Control, Data access and Communication (CODAC)
- Cooling water
- Cryogenics
- Diagnostics
- Fuel cycle
- Hot cell a secure environment for processing, repair or testing, etc., of components that have become activated by neutrons.
- Power supply
- Remote handling
- Heating and current drive
- Vacuum system


ITER is being assembled






ITER is being assembled





There is a long way to go, but we are on the right path...






Dec 2025 First Plasma

2035

Deuterium-Tritium Operation begins

Joint European Torus (JET) facility has a recordbreaking 59 megajoules of sustained fusion energy







 Record-breaking 59 megajoules of sustained fusion energy in Joint European Torus (JET) facility in Oxford demonstrates powerplant potential and strengthens case for ITER.

Stellarator uses twisted coil to generate poloidal magnetic field






Stellarator



A figure-8 stellarator solved the drift issues





A figure-8 stellarator solved the drift issues



Lyman Spitzer, Jr. came out the idea during a long ride on a ski lift at Garmisch-Partenkirchen





Model A stellarator








Exhibit model of a figure-8 stellarator for the Atoms for Peace conference in Geneva in 1958





Twisted magnetic field lines can be provided by toroidal coils with helical coils






Construction of a pair of helical magnetic coils for the Advanced Toroidal Facility torsatron







LHD stellarator in Japan (Heliotron)











https://en.wikipedia.org/wiki/Compact_Toroidal_Hybrid https://www.energyencyclopedia.com/en/glossary/heliotron https://en.wikipedia.org/wiki/Large_Helical_Device


Wendelstein 7-X is a stellarator built by Max Planck Institute for Plasma Physics (IPP)



Demonstration of a magnetic mirror machine








Show video.


Plasma is partially confined by the magnetic field







Many mirror points are provided by a pair of ring-type magnets










Outline



- Introduction to nuclear fusion
- Magnetic confinement fusion (MCF)
 - Tokamak
 - Stellarator
- Inertial confinement fusion (ICF)
 - Indirection drive ICF
 - Direct drive ICF
- Innovation idea MCF + ICF
- Plasma in space
- Pulsed-power system at NCKU


Compression happens when outer layer of the target is heated by laser and ablated outward




Inertial confinement fusion: an introduction, Laboratory for Laser Energetics, University of Rochester R. Betti, HEDSA HEDP Summer School, 2015

Plasma is confined by its own inertia in inertial confinement fusion (ICF)





Spatial profile at stagnation




A ball can not be compressed uniformly by being squeezed between several fingers






Rayleigh-Taylor instability



Stages of a target implosion





P.-Y. Chang, PhD Thesis, U of Rochester (2013)

R. S. Craxton, etc., *Phys. Plasmas* 22, 110501 (2015)

A spherical capsule can be imploded through directly or indirectly laser illumination







Rochester is known as "The World's Image Center"





There are many famous optical companies at Rochester



Kodak





Eastman school of music


BAUSCH+LOMB



Laboratory for Laser Energetics, University of Rochester is a pioneer in laser fusion

- OMEGA Laser System
 - 60 beams
 - >30 kJ UV on target
 - 1%~2% irradiation nonuniformity
 - Flexible pulse shaping












 Can propagate to the OMEGA or OMEGA EP target chamber








The OMEGA Facility is carrying out ICF experiments using a full suite of target diagnostics



The 1.8-MJ National Ignition Facility (NIF) will demonstrate ICF ignition and modest energy gain






OMEGA experiments are integral to an ignition demonstration on the NIF.


Targets used in ICF





Triple-point temperature : 19.79 K






http://www.lle.rochester.ed https://en.wikipedia.org/wiki/Inertial_confinement_fusion R. S. Craxton, etc., *Phys. Plasmas* **22**, 110501 (2015)

Targets used in ICF







Cryogenic shroud



a Cryogenic hohlraum




Rugby hohlraum



b



d Tent holder



https://www.lle.rochester.edu/index.php/2014/11/10/next-generation-cryo-target/ Introduction to Plasma Physics and Controlled Fusion 3rd Edition, by Francis F. Chen https://www.llnl.gov/news/nif-shot-lights-way-new-fusion-ignition-phase

Softer material can be compressed to higher density

Compression of a baseball



Compression of a tennis ball









A shock is formed due to the increasing sound speed of a compressed gas/plasma








Acoustic/compression wave driven by a piston:




Nature letter "Fuel gain exceeding unity in an inertially confined fusion implosion"





Fuel gain exceeding unity was demonstrated for the first time.

The hot spot has entered the burning plasma regime




2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021

Year

A. L. Kritcher, etc., Phys. Plasmas, 28, 072706 (2021)

H. F. Robey, etc., Phys. Plasmas, 25, 012711 (2018)


National Ignition Facility (NIF) achieved a yield of more than 1.3 MJ from ~1.9 MJ of laser energy in 2021 (Q~0.7)



National Ignition Facility (NIF)
 achieved a yield of more than 1.3
 MJ (Q~0.7). This advancement puts
 researchers at the threshold of
 fusion ignition.

THE ROAD TO IGNITION

The National Ignition Facility (NIF) struggled for years before achieving a high-yield fusion reaction (considered ignition, by some measures) in 2021. Repeat experiments, however, produced less than half the energy of that result.




 Laser-fusion facility heads back to the drawing board.

T. Ma, ARPA-E workshop, April 26, 2022

J. Tollefson, Nature (News) 608, 20 (2022)


"Ignition" (target yield larger than one) was achieved in NIF on 2022/12/5




https://physicstoday.scitation.org/do/10.1063/PT.6.2.20221213a/full/ The age of ignition: anniversary edition, LLNL-BR-857901

A gain over 4 has been achieved








 In recent attempts, the team at NIF increased the yield of the experiment, first to 5.2 megajoules and then again to 8.6 megajoules, according to a source with knowledge of the experiment.

External "spark" can be used for ignition




Shock ignition




Fast ignition

a) channeling FI concept




b) cone-in-shell FI concept




A shock is formed due to the increasing sound speed of a compressed gas/plasma








Acoustic/compression wave driven by a piston:




Ignition can happen by itself or being triggered externally




Self-ignition



External "spark" for fast ignition






Outline



- Introduction to nuclear fusion
- Magnetic confinement fusion (MCF)
 - Tokamak
 - Stellarator
- Inertial confinement fusion (ICF)
 - Indirection drive ICF
 - Direct drive ICF
- Innovation idea MCF + ICF
- Plasma in space
- Pulsed-power system at NCKU

A strong magnetic field reduces the heat flux





$$oldsymbol{q}_{ au} = -\kappa_{||}
abla_{||} oldsymbol{T} - \kappa_{\perp}
abla_{\perp} oldsymbol{T}$$
 $oldsymbol{\kappa}_{||} = \kappa_{0} oldsymbol{T}^{5/2}$

$$\kappa_{||} = \kappa_0 T^{5/2}$$

$$\kappa_{\perp} = \frac{\kappa_{||}}{\chi^2}$$
 for large Hall parameter $\chi \propto \frac{I_{\text{mfp}}}{R_{\perp}} >> 1$

Typical hot spot conditions:


 $R_{hs} \sim 40 \ \mu m, \ \rho \sim 20 \ g/cm^3, \ T \sim 5 \ keV$:

 $B > 10 \, \text{MG}$ is neededfor $\chi > 1$

Magnetic-flux compression can be used to provide the needed magnetic field.

Principle of frozen magnetic flux in a good conductor is used to compress fields






$$\Phi = \pi r_0^2 B_0 = \pi r^2 B$$

Plasma can be pinched by parallel propagating plasmas







Plasma can be heated via pinches








Theta pinch



Sandia's Z machine is the world's most powerful and efficient laboratory radiation source





Stored energy: 20 MJ

Marx charge voltage: 85 kV

Peak electrical power: 85 TW

Peak current: 26 MA

Rise time: 100 ns


Peak X-ray emissions: 350 TW

Peak X-ray output: 2.7 MJ

Z machine








Z machine







Stored energy: 20 MJ

Peak electrical power: 85 TW


Peak current: 26 MA

Rise time: 100 ns

Peak X-ray output: 2.7 MJ

Z machine discharge






Before and after shots

Before shots




After shots





SAND2017-0900PE_The sandia z machine - an overview of the world's most powerful pulsed power facility.pdf

Promising results were shown in MagLIF concept conducted at the Sandia National Laboratories





The stagnation plasma reached fusion-relevant temperatures with a 70 km/s implosion velocity


S. A. Slutz *et al* Phys. Plasmas 17 056303 (2010) M. R. Gomez *et al* Phys. Rev. Lett. 113 155003 (2014)

MagLIF target











Neutron yield increased by 100x with preheat and external magnetic field.










Sheared flow stabilizes MHD instabilities











$$m = 1 (kink)$$













$$\frac{dV_Z}{dr}\neq 0$$


- M. G. Haines, etc., Phys. Plasmas 7, 1672 (2000)
- U. Shumlak, etc., Physical Rev. Lett. 75, 3285 (1995)
- U. Shumlak, etc., ALPHA Annual Review Meeting 2017

A z-pinch plasma can be stabilized by sheared flows










https://www.zapenergyinc.com/about


A. D. Stepanov, etc., Phys. Plasmas 27, 112503 (2020)


Elevated electron temperature coincident with observed fusion reactions in a sheared-flow-stabilized z pinch



B. Levitt, etc., Phys. Rev. Lett. **132**, 155101 (2024)

Fusion reactor concept by ZAP energy






https://www.zapenergyinc.com/about E. G. Forbes, etc., Fusion Sci. Tech. 75, 599 (2019)


There are alternative






Commonwealth Fusion Systems, a MIT spin-out company, is building a high-magnetic field tokamak





- The fusion gain Q > 2 is expected for SPARC tokamak.

Merging compression is used to heat the tokamak at the start-up process in ST40 Tokamak at Tokamak Energy Ltd




- High temperature superconductors are used.
- $B_T \sim 3 T$



M. Gryaznevich, etc., Fusion Eng. Design, **123**,177 (2017) https://www.tokamakenergy.co.uk/


Reconnection





https://www.youtube.com/watch?v=7sS3Lpzh0Zw

Merging compression is used to heat the plasma




http://www.100milliondegrees.com/merging-compression/P. F. Buxton, etc., Fusion Eng. Design, **123**, 551 (2017)


Spherical torus (ST) and compact torus (CT)




Spherical torus (ST)



- Compact torus (CT)
 - Spheromak




Field reversed configuration (FRC)




Zhe Gao, Matter Radiat. Extremes **1**, 153 (2016) https://en.wikipedia.org/wiki/Field-reversed_configuration

Field reverse configuration is used in Tri-alpha energy








^{*}Magneto-Inertial Fusion & Magnetized HED Physics by Bruno S. Bauer, UNR & Magneto-Inertial Fusion Community

^{**}https://en.wikipedia.org/wiki/Field-reversed configuration


Field reverse configuration is used in Tri-alpha energy





NBI for Tri-Alpha Energy Technologies





Neutral beams are injected in to the chamber for spinning the FRC






https://tae.com/media/


FRC sustain longer with neutral beam injection






General fusion is a design ready to be migrated to a power plant








A spherical tokamak is first generated





Plasma injector for the spherical tokamak









A spherical tokamak is generated in a liquid metal vortex





The spherical tokamak is compressed by the pressure provided by the sournding hydraulic pistons





BBC: General Fusion to build its Fusion Demonstration Plant in the UK, at the UKAEA Culham Campus

Nuclear energy: Fusion plant backed by Jeff Bezos to be built in UK

By Matt McGrath Environment correspondent

(1) 17 June






A company backed by Amazon's Jeff Bezos is set to build a large-scale nuclear fusion demonstration plant in Oxfordshire.

Canada's General Fusion is one of the leading private firms aiming to turn the

Helion energy is compressing the two merging FRCs






https://www.helionenergy.com/

Two FRCs are accelerated toward each other






Two FRCs merge with each other





The merged FRC is compressed electrically to high temperature







Similar concept will be studied in our laboratory.

Projectile Fusion is being established at First Light **Fusion Ltd, UK**







 I_{peak} =14 MA w/ T_{rise} ~2us.









High pressure is generated by the colliding shock.

A gas gun is used to eject the projectile









https://www.youtube.com/watch?v=JN7lyxC11n0 https://www.youtube.com/watch?v=aW4eufacf-8

Many groups aim to achieve ignition in the MCF regime in the near future


ITER – 2025 First Plasma
 2035 D-T Exps
 2050 DEMO



- Tokamak energy, UK
 - 2025 Gain
 - 2030 to power grid



Commonwealth Fusion Systems, USA – 2025 Gain



https://www.iter.org

https://www.tokamakenergy.co.uk/ https://www.psfc.mit.edu/sparc

Fusion is blooming

































































































We are closed to ignition!




We are closed to ignition!




We are closed to ignition!





A. J. Webster, Phys. Educ. **38**, 135 (2003) R. Betti, etc., Phys. Plasmas, **17**, 058102 (2010)

Formosa Integrated Research Spherical Tokamak (FIRST) aiming for the first plasma in 2026




- *R/a/b*=45/32/76.8 cm
- Aspect ratio = R/a = 1.5
- k = b/a = 2.4
- $\delta = 0.5$
- $B_T = 0.1 \sim 0.5 \text{ T}$
- T≥ 100 eV
- $I_p \ge 100 \text{ kA}$
- Gas: H₂
- Ohmic heating
- Duration: 100 ms
- FIRST is targeted for
 - Low aspect ratio
 - High beta
 - High bootstrap current
- Short term goal in my group study the startup mechanism.

We welcome anyone interested in fusion research to join us!


Temperature of 100 eV is the threshold of radiation barrier by impurities





This is a joint project including 6 groups

- 馬維揚團隊 @ 國家原子能科技研究院
 - Site
- 張博宇團隊 @ 成功大學 太空與電漿科學研究所
 - System design and development/diagnostics
- 向克強團隊 @ 成功大學 前瞻電漿中心
 - Theoretical design
- 河森榮一郎團隊 @ 成功大學 太空與電漿科學研究所
 - Diagnostics
- 柳克強團隊 @ 清華大學 工程與系統科學系
 - Diagnostics
- 蔡宗哲團隊 @ 國家高速網路與計算中心
 - Simulation
- 張存續團隊 @ 清華大學 物理系
 - RF startup
 - We welcome anyone interested in fusion research to join us!



https://www.sipa.gov.tw/home.jsp?mserno=20100 1210037&serno=201001210041&menudata=Chin eseMenu&contlink=content/introduction_4_1.jsp&serno3=201002010023

Students are encouraged to study either science or engineering



- Management team @ Laboratory for Laser Energetics, University of Rochester
 - Administrative \ Engineering \ Experimental \ Omega Laser Facility \ Plasma & Ultrafast Laser Science & Engineering \ Theory division \ Laboratory Safety \ Laser and Materials Technology
- Science involved:
 - Plasma Physics ` Physics ` Material Science ` Super conductors ` Chemistry ` Mathematics...
- Engineering involved:
 - Electrical Engineering \ Mechanical Engineering \ Computer science \ Artificial intelligent \ Control \ Optics \ ...

Skip 223

Outline



- Introduction to nuclear fusion
- Magnetic confinement fusion (MCF)
 - Tokamak
 - Stellarator
- Inertial confinement fusion (ICF)
 - Indirection drive ICF
 - Direct drive ICF
- Innovation idea MCF + ICF
- Plasma in space
- Pulsed-power system at NCKU

Aurora






https://en.wiktionary.org/wiki/aurora

Aurora seen from a satellite







https://flashpack.com/insights/2014/11/20/aurora-australis-forget-the-northern-lights-have-you-heard-about-the-southern-lights/

Earth's magnetic field








https://www.nasa.gov/mission_pages/sunearth/news/gallery/Earths-magneticfieldlines-dipole.html http://www.pas.rochester.edu/~blackman/ast104/emagnetic.html


Earth magnetic fields are strongly influenced by solar wind





Reconnection







https://www.youtube.com/watch?v=7sS3Lpzh0Zw

Corona mass ejection (CME)








http://cse.ssl.berkeley.edu/SegwayEd/lessons/exploring_magnetism/in_Solar_Flares/s4.html#sf

Reconnections occur in many locations






The Aurora Borealis:

https://www.youtube.com/watch?v=IT3J6a9p_o8


Planeterrella is an aurora simulator






Simple glow discharge is demonstrated





Aurora/ring current are demonstrated






- B w/ magnet: aurora demonstration
- F w/ magnet: ring current

Aurora and ring current are expected to be seen







Neutral beam source



- Neutral beam injection for heating plasma in Tokamak
 - Jure Maglica, Seminar at University in Ljubljana
 - lan G. Brown, The Physics and Technology of Ion Sources
- Electric propulsion (plasma thrusters)
 - D. M. Goebel and I. Katz, Fundamentals of Electric Propulsion: Ion and Hall Thrusters


Neutral beam source




- Neutral beam injection for heating plasma in Tokamak
 - Jure Maglica, Seminar at University in Ljubljana
 - Ian G. Brown, The Physics and Technology of Ion Sources
- Electric propulsion (plasma thrusters)
 - D. M. Goebel and I. Katz, Fundamentals of Electric Propulsion: Ion and Hall Thrusters

Hot plasma is confined by the magnetic field in magnetic confinement fusion








Neutral beam injector is one of the main heat mechanisms in MCF





Varies way of heating a MCF device




Sy	rstem	Frequency/ energy	Maximum power coupled to plasma	Overall system efficiency	Development/ demonstration required	Remarks
ECRF	Demonstrated in tokamaks	28–157 GHz	2.8 MW, 0.2 s	30-40%	Power sources and windows, off-axis CD	Provides off-axis CD
	ITER needs	150–170 GHz	50 MW, SS			
ICRF	Demonstrated in tokamaks	25–120 MHz	22 MW, 3 s (L-mode); 16.5 MW, 3 s (H-mode)	50-60%	ELM tolerant system	Provides ion heating and smaller ELMs
	ITER needs	40–75 MHz	50 MW, SS			
LHRF	Demonstrated in tokamaks	1.3–8 GHz	2.5 MW, 120 s; 10 MW, 0.5 s	45–55%	Launcher, coupling to H-mode	Provides off-axis CD
	ITER needs	$5~\mathrm{GHz}$	50 MW, SS			
+ve ion NBI -ve ion	Demonstrated in tokamaks	$80140~\mathrm{keV}$	40 MW, 2 s; 20 MW, 8 s	35-45%	None	Not applicable
	ITER needs	None	None			
	Demonstrated in tokamaks	$0.35\;\mathrm{MeV}$	$5.2 \mathrm{MW}, \mathrm{D}^-, 0.8 \mathrm{s}$ (from 2 sources)			
	ITER needs	$1~{ m MeV}$	50 MW, SS	~37%	System, tests on tokamak, plasma CD	provides rotation

^{&#}x27;S S' indicates steady state

Neutral particles heat the plasma via coulomb collisions






- 1. create energetic (fast) neutral ions
- 2. ionize the neutral particles
- 3. heat the plasma (electrons and ions) via Coulomb collisions

Negative ion source is preferred due to higher neutralization efficiency






There are two ways to make negative ions – surface and volume production



- Surface production, depends on :
 - Work function Φ





- Perpendicular velocity
- Work function can be reduced by covering the metal surface with cesium

$$H + e^- \rightarrow H^-$$


 $H^+ + 2e^- \rightarrow H^-$

Volume production:

$$H_2 + e_{\textit{fast}}(>20 \text{ eV}) \rightarrow H_2^*(\text{excited state}) + e_{\textit{fast}},$$
 $H_2^*(\text{excited state}) + e_{\textit{slow}}(\approx 1 \text{ eV}) \rightarrow H^- + H.$


Two-chamber method of negative ions in volume production with a magnetic filter






Adding cesium increases negative ion current








Electrons need to be filtered out since they are extracted together with negative ions






Acceleration



Multi-stage acceleration




The ITER neutral beam system: status of the project and review of the main technological issues, presented by V. Antoni

NBI system of the LHD fusion machine








INSULATOR

- 180 keV and 30 A
- Arc chamber: 35 cm x 145 cm, 21cm in depth
- Single stage accelerator

JT60U NBI system



- JT-60 (Japan-Torus) is a tokamak in Japan.
- 550 keV, 22A
- 2m in diameter and 1.7 m in height
- 3-stage accelerator



Neutralization



Gas neutralization

Collisions between fast negative ions and atoms

$$H^- + H_2 \longrightarrow H + H_2 + e^-$$

Fast ions can lose another electron after neutralized

$$H + H_2 \rightarrow H^+ + H_2 + e^-$$


- Plasma neutralization
 - Collisions with charged particles in plasma

$$H^{-} + X(e, Ar, H^{+}, H_{2}^{+}) \longrightarrow H + X + e^{-}$$

- The efficiencies reach up to 85% for fully ionized hydrogen plasma

Beam dump






NBI for ITER




- beam components (Ion Source, Accelerator, Neutralizer, Residual Ion Dump and Calorimeter)
- other components (cryo-pump, vessels, fast shutter, duct, magnetic shielding, and residual magnetic field compensating coils)



The ITER neutral beam system: status of the project and review of the main technological issues, presented by V. Antoni

Neutral beam penetration






- Parallel direction
 - Longest path through the densest part of the plasma
 - Harder to be built
- Perpendicular direction
 - Path is short
 - Larger perpendicular energies leads to larger losses
 - Easier to be built

NBI for Tri-Alpha Energy Technologies





Neutral beams are injected in to the chamber for spinning the FRC





Outline



- Introduction to nuclear fusion
- Magnetic confinement fusion (MCF)
 - Tokamak
 - Stellarator
- Inertial confinement fusion (ICF)
 - Indirection drive ICF
 - Direct drive ICF
- Innovation idea MCF + ICF
- Plasma in space
- Pulsed-power system at NCKU

Neutral beam source




- Neutral beam injection for heating plasma in Tokamak
 - Jure Maglica, Seminar at University in Ljubljana
 - lan G. Brown, The Physics and Technology of Ion Sources
- Electric propulsion (plasma thrusters)
 - D. M. Goebel and I. Katz, Fundamentals of Electric Propulsion: Ion and Hall Thrusters


Comparison between liquid rockets and ion thrusters



Liquid rockests

- u~4500 m/s
- Isp~450 s
- Energy ~ 100GJ
- Power ~ 300MW
- Thrust ~ 2x10⁶ N
- Ion thrusters
 - u~30000 m/s
 - Isp~3000 s
 - Energy ~ 1000GJ
 - Power ~ 1kW
 - Thrust ~ 0.1 N






Electric thruster types - electrothermal



Resistojet

Resistojet AC or DC Power Hot Gas Exhaust Hydrogen, Ammonia, or Decomposed Hydrazine 300 s (hydrazine) 900 s (hydrogen)

Arcjet





Electric thruster types - electrostatic



Ion thruster









Electric thruster types - Electromagnetic



Pulsed plasma thruster



 Magnetoplasmadynamic thruster (MPD)



The thrust in an ion engine is transferred by the electrostatic force between the ions and the two grids



$$\frac{dE(x)}{dx} = \frac{\rho(x)}{\varepsilon_0} = \frac{qn_i(x)}{\varepsilon_0}$$


$$E(x) = \frac{q}{\varepsilon_0} \int_0^x n_i(x') dx' + E_{\text{screen}}$$

Gauss's law: $\sigma = \varepsilon_0 E_{\text{screen}}$

$$F_{\text{screen}} = \sigma \frac{(E_{\text{screen}} + 0)}{2} = \frac{1}{2} \varepsilon_0 E_{\text{screen}}^2$$

$$F_{\text{accel}} = -\sigma \frac{(E_{\text{accel}} + 0)}{2} = -\frac{1}{2} \varepsilon_0 E_{\text{accel}}^2$$

$$T = F_{\text{screen}} + F_{\text{accel}} = \frac{1}{2} \varepsilon_0 (E_{\text{screen}}^2 - E_{\text{accel}}^2)$$



$$F_{\text{ion}} = q \int_0^d n_i(x) E(x) dx = \varepsilon_0 \int_0^d \frac{dE}{dx} E dx = \frac{1}{2} \varepsilon_0 (E_{\text{accel}}^2 - E_{\text{screen}}^2)$$

The rocket equation



$$p(t) = p(t + dt)$$

$$Mv = (M - dm_p)(v + dv) + dm_p(v - v_{ex})$$

$$Mv = Mv + Mdv - dm_pv - dm_pdv + dm_pv - dm_pv_{ex}$$


 $dv \sim -v_{\rm ex} rac{dM}{M}$ where $dm_{\rm p} dv$ is neglected and $dm_{\rm p} = -dM$


$$\int_{v_i}^{v_f} dv = -v_{\text{ex}} \int_{m_d + m_p}^{m_d} \frac{dM}{M}$$

$$\Delta \mathbf{v} = (\mathbf{Isp} \times \mathbf{g}) \ln \left(\frac{\mathbf{m}_d + \mathbf{m}_p}{\mathbf{m}_d} \right)$$

$$egin{aligned} m{m}_p &= m{m}_d [e^{\Delta ext{v}/v_{ ext{ex}}} - m{1}] \ &= m{m}_d [e^{\Delta ext{v}/(ext{Isp} imes g)} - m{1}] \end{aligned}$$







Force transfer



$$T = -\frac{d}{dt}(m_p v_{\text{ex}}) = -v_{\text{ex}} \frac{dm_p}{dt} = -\dot{m}_p v_{ex}$$

$$\dot{m}_p = QM$$

$$P_{
m jet} = -rac{1}{2}\dot{m}_p v_{
m ex}^2 = -rac{T^2}{2\dot{m}_p}$$
 $Q = ext{propellant particle}$ $M = ext{atomic mass in kg}$

$$T = -\frac{dm_{\rm p}}{dt}v_{\rm ex} \approx -\dot{m}_i v_i$$

$$v_i = \sqrt{\frac{2qV_b}{M}}$$

$$-\dot{m}_i = \frac{I_b M}{q}$$

 m_p = propellant mass flow rate in kg/s

Q = propellant particle flow rate in particles/s

 $\dot{m}_i = \text{ion mass flow rate in kg/}s$ $I_b = \text{ion current}$

$$T = \sqrt{\frac{2M}{e}} I_b \sqrt{V_b} \text{ (Nt)}$$


Ion thruster has the highest specific impulse (Isp)




Thruster	Specific Impulse (s)	Input Power (kW)	Efficiency Range (%)	Propellant
Cold gas	50-75			Various
Chemical (monopropellant)	150-225	—		N_2H_4 H_2O_2
Chemical (bipropellant)	300-450		_	Various
Resistojet	300	0.5-1	65-90	N ₂ H ₄ monoprop
Arcjet	500-600	0.9-2.2	25-45	N ₂ H ₄ monoprop
Ion thruster	2500-3600	0.4-4.3	40-80	Xenon
Hall thrusters	1500-2000	1.5-4.5	35-60	Xenon
PPTs	850-1200	<0.2	7–13	Teflon

Metallic Ion Thruster Using Magnetron E-Beam Bombardment (MIT-MEB)





Electrons are used to generate metallic gas, metallic plasma and to neutralize ions





High energy particle accelerator



- linear particle accelerator (Linac)
- Cyclotron

Synchrotron









 Reference: Introduction to plasma phenomena and plasma medicine, Y. Nishida and K.-L. Ou

A linear particle accelerator (linac) accelerates charged particles using a series of oscillating electric potentials along a linear beamline



Cyclotrons use a magnetic field to cause particles to move in circular orbits





$$\omega_{\rm ce} = \frac{eB}{m_e c}$$


$$r_e = rac{v}{\omega_{
m ce}} = rac{m_e c v}{e B}$$

Cyclotron was invented by Ernest Lawrence who earned the 1939 Nobel price in physics

270


Synchrotron uses time-dependent guiding magnetic field synchronized to a particle beam





Stanford linear accelerator center (SLAC) is a 50 GeV electron / positron accelerator





Large Hadron Collider (LHC) is the world's largest and most powerful particle collider providing 13 TeV protons



Plasma based accelerators will become 3 orders smaller than the regular microwave based accelerator



- Maximum field strength:
 - Microwave: 100 MV/m
 - Plasma: >10 GV/m, 300 GV/m was achieved using laser wakefield accelerator¹
- Plasma based high energy accelerators:
 - V_pxB or surfatron accelerator²
 - Plasma wakefield accelerator (PWFA)³
 - Plasma beat wave accelerator (PBWA)⁴
 - Laser wakefield accelerator (LWFA)⁴

¹N. A. M. Hafz, *et al.*, Nature Photonics **2**, 571 (2008)


²T. Katsouleas and J. Dawson, Phys. Rev. Lett. **51**, 392 (1983)


³P. Chen, et al., Phys. Rev. Lett. **54**, 693 (1985)

⁴T. Tajima and J. M. Dawson, Phys. Rev. Lett. **43**, 267 (1979)

Charged particles can be accelerated in the wave electric field







Who will catch the wave?





Plasma wake field accelerator is just like boat wake surfing







A wake surfer catches the wake field via being pulled by the boat using a roap







https://www.youtube.com/watch?v=VFp7SloeAnk


The surfer glides in a direction not parallel to the wave direction to be in phase to the wave propagation






The surfer glides in a direction not parallel to the wave direction to be in phase to the wave propagation





Electrons may be accelerated to speed of light using $V_p x B$ acceleration (Surfatron)





 Plane wave electric field and uniform magnetic field:

$$\vec{E} = E_0 \sin(kx - \omega t)\hat{x}$$

$$\vec{B} = B\hat{z}$$

$$\frac{d}{dt}(\gamma v_x) = \frac{qE_0}{m} \sin(kx - \omega t) + \omega_c v_y$$


$$\frac{d}{dt}(\gamma v_y) = -\omega_c v_x$$

$$\gamma = \frac{1}{1 - \frac{v_x^2 + v_y^2}{c^2}}$$

On the wave frame and if the particle is trapped in the wave:

$$x_1 = x - v_{\text{ph}}t$$
 $\frac{d}{dt}(\gamma v_{\chi}) = 0$
 $v_{\chi} \rightarrow v_{\text{ph}}$

$$v_y = -\frac{\omega_c v_{\rm ph} t}{\gamma_{\rm ph} \sqrt{1 + \frac{\omega_c^2 t^2 v_{\rm ph}^2}{c^2}}}$$




[•] T. Katsouleas, et al., PRL 51, 392 (1983)

[•] T. Katsouleas, et al., IEEE TNS. NS-30, 3241 (1983)

Experimental results of V_pxB acceleration (Surfatron)







- $n_0 \sim 1-30 \times 10^{17} \text{ m}^{-3}$
- $T_e \sim 2-5 \text{ eV}$

- T_i ~ 0.1-0.2 eV
- Microwave frequency: 3-10 GHz

Plasma beat wave accelerator







$$sin(x_1) + sin(x_2) = 2 sin\left(\frac{x_1 + x_2}{2}\right) cos\left(\frac{x_1 - x_2}{2}\right)$$

A plasma wave is driven by the laser beat wave




$$\omega_0 = \omega_2 - \omega_1$$

$$k_0 = k_2 - k_1$$

$$v_{\mathrm{ph}} = v_g = c \sqrt{1 - \frac{{\omega_p}^2}{{\omega_0}^2}}$$


$$F = -e\nabla \phi_p = -\nabla \frac{e^2 E^{(1)} \cdot E^{(2)*}}{m\omega_1\omega_2}$$

Plasma wave



Electrons were accelerated to over 20 MeV using plasma beat wave accelerator








Intensity: $2x10^{13} \sim 2x10^{14} \text{ W/cm}^2$

Injected E-beam: 0.1~1 MeV


 $n_0 = 3x10^{16} \sim 7x10^{17} \text{ cm}^{-3}$




- O 246J/4e17/cc

Plasma wakefield accelerator employs two beams


- When a bunch of electrons enter the plasma, they expel local electrons.
- When the bunch of electrons leave the plasma, the local electrons try to return but oscillate around their original locations and generate a wake field behind the bunch.
- The longitudinal field of the wake can accelerate the particles in the back.
- Key components:
 - Drive bunch: excite wakefield
 - Test bunch: beam that is accelerated to high energy





Energy doubling of 42 GeV electrons in a metre-scale plasma wakefield accelerator





Dream beam – the dawn of compact particle accelerators





Ponderomotive force expelled electrons away from the higher electric field region



$$m_s\ddot{x}=q_sE=q_sE_0(x)\cos\omega t$$
 $x=x_0+x_1$ where $x_0=\overline{x}$ $m_s(\ddot{x}_0+\ddot{x}_1)=q_s\left(E_0+x_1rac{\mathrm{d}E_0}{\mathrm{d}x}
ight)\cos\omega t$

Take time average:

$$m_s \ddot{x}_0 = q_s \frac{dE_0}{dx} \bigg|_{x_0} \overline{x_1 \cos \omega t}$$
• $\ddot{x}_1 \gg \ddot{x}_0$, $E_0 >> x_1 \frac{dE_0}{dx}$
 $m_s \ddot{x}_1 = q_s E_0 \cos \omega t$

$$x_1 = -\frac{q_s E_0}{m_s \omega^2} \cos \omega t$$

$$\ddot{x}_0 = -\frac{q_s^2 E_0}{2m_s^2 \omega^2} \frac{dE_0}{dx}$$

$$\frac{dE_0}{dx} = 0 \qquad \frac{dE_0}{dx} > 0$$
Weak Strong


$$F_p = m_s \ddot{x}_0 = -\frac{q_s^2}{4m_s \omega^2} \frac{d}{dx} (E_0^2)$$

Laser is used to create a bunch in laser wakefield accelerator




$$I(r,z) = \frac{2P}{\pi w^{2}(z)} \exp \left[-\frac{2r^{2}}{w^{2}(z)}\right]$$

- Waist: $w(z) = w_0 \sqrt{1 + \frac{z^2}{{z_R}^2}}$
- Rayleigh length: $z_R = \frac{\pi w_0^2}{\lambda_L}$




Bubble/blow-out regime






A plasma wake is generated by a short pulse laser








V. Malka, *et al.*, Nature Physics **4**, 447 (2008) http://cuos.engin.umich.edu/researchgroups/hfs/research/laser-wakefield-acceleration/


The wakefield generated by a short pulse laser is very similar to the wave behind a boat





Ionization injection






- Large relative energy spread
- Energy required to trap electrons is reduced so that electron beams with large charge can be produced in a moderate laser energy


Colliding laser pulses injection






Few femtosecond, few kiloampere electron bunch is produced by a laser-plasma accelerator







Plasma medicine



Reference:

- "Applied Plasma Medicine", by G. Fridman, et al., Plasma Process.
 Polym., 5, 503, 2008
- "Plasma Medicine", by A. Fridman and G. Fridman

Outline



- Example of several plasma discharges for plasma medicine
- Living tissue sterilization
- Blood coagulation
- Nitrogen oxide (NO) treatment
- Non-thermal plasma treatment of melanoma skin cancer
- Skin regeneration
- Egg sterilization
- Facemask regeneration

Outline



- Example of several plasma discharges for plasma medicine
- Living tissue sterilization
- Blood coagulation
- Nitrogen oxide (NO) treatment
- Non-thermal plasma treatment of melanoma skin cancer
- Skin regeneration
- Egg sterilization
- Facemask regeneration

Plasma is characterized by the electron and ion temperatures



- Non-thermal plasma
 - $-T_i \ll T_e$
 - Also called non-equilibrium plasma
- Thermal plasma
 - T_i≈T_e
- Earlier applications of plasma in medicine thermal effects of plasma


Plasma can provide good surface treatment with low temperature





Treatment	Surface treatment level	Depth	Temperature	Cost
Chemical	Large	Deep	Room temperature ~200 °C	Medium
Heat	Only oxidizing	Deep	High temperature	Cheap
Radiation	Small	Whole sample	High temperature	Expensive
Plasma	Large	Surface	Room temperature ~100 °C	Cheap ~ Medium

Microwave plasma torch











Dielectric-barrier discharges (DBDs)









Plasma-needle discharge






Atmospheric-pressure cold helium microplasma jets





Space charge effect enhance the electric field







J. L. Walsh, et al., J. Phys. D: Appl. Phys., 43, 075201 (2010)

There are three different modes: chaotic, bullet, and continuous mode








In bullet mode, the plasma jet comes out as a pulse




 wavelength-integrated optical emission signal (350–800 nm) Images of bullet mode



Floating-electrode dielectric barrier discharge (FE-DBD)








Simplified electrical schematic of FE-DBD



- electrode itself
- electrode near the treated object
- e-plasma discharge








$$C_{\rm ob} >> C_{\rm gap} \Rightarrow V_{\rm ob} << V_{\rm gap}$$


Depending on the needs, the size and the shape of FE-DBD treatment electrodes can vary





G. Fridman, et al., Plasma Chem. Plasma Process., 26, 425 (2006)

FE-DBD is a direct plasma medicine



G. Fridman, *et al.*, Plasma Chem. Plasma Process., **26**, 425 (2006) Plasma medicine, by Alexander Fridman and Gary Friedman

Outline



- Example of several plasma discharges for plasma medicine
- Living tissue sterilization
- Blood coagulation
- Nitrogen oxide (NO) treatment
- Non-thermal plasma treatment of melanoma skin cancer
- Skin regeneration
- Egg sterilization
- Facemask regeneration

Bacteria concentration reduces after being treated with FE-DBD



Table 1. Bacteria sterilization results (in cfu⋅mL⁻¹).^[26]

Original concentration	5 s of FE-DBD	10 s of FE-DBD	15 s of FE-DBD
10 ⁹	850 ± 183	9 ± 3	4 ± 4
10 ⁸	22 ± 5	5 ± 5	0 ± 0
10 ⁷	6 ± 6	0 ± 0	0 ± 0

 Maximum acceptable dose – the highest dose that doesn't cause a damage on skin

The power of FE-DBD is low enough such that the tissue is not damaged by the plasma







Figure 6.23 Photos (top) and tissue histology (bottom) of cadaver skin samples after FE-DBD treatment: (a, d) control; (b, e) after 15 s of treatment; and (c, f) after 5 min of treatment – no visible damage is detected.

G. Fridman, *et al.*, Plasma Chem. Plasma Process., **26**, 425 (2006) Plasma medicine, by Alexander Fridman and Gary Friedman

Bacteria is inactivated by the plasma



- ~1.3x10⁷ cfu/cm² (10⁹ cfu/ml) of skin flora
 (CFU: colony-forming unit)
- Treated by FE-DBD plasma for 10 s




Outline




- Example of several plasma discharges for plasma medicine
- Living tissue sterilization
- Blood coagulation
- Nitrogen oxide (NO) treatment
- Non-thermal plasma treatment of melanoma skin cancer
- Skin regeneration
- Egg sterilization
- Facemask regeneration


Plasma can stimulate blood coagulation












G. Fridman, et al., Plasma Chem. Plasma Process., 26, 425 (2006)

Example of blood coagulation using plasma







Saphenous vein is a major blood vessel for a mouse

(a)



If left untreated following a cut animal will bleed out (control)

(b)

15 seconds at 0.8 Watt/cm² stops the bleeding completely right after treatment

-)

G. Fridman, *et al.*, Plasma Process. Polym., **5**, 503 (2008) G. Fridman, *et al.*, Plasma Chem. Plasma Process., **26**, 425 (2006) Plasma medicine, by Alexander Fridman and Gary Friedman

Outline



- Example of several plasma discharges for plasma medicine
- Living tissue sterilization
- Blood coagulation
- Nitrogen oxide (NO) treatment
- Non-thermal plasma treatment of melanoma skin cancer
- Skin regeneration
- Egg sterilization
- Facemask regeneration

Nitrogen oxide (NO) serves a multitude of essential biological functions



- Blood coagulation
- Immune system
- Early apoptosis (細胞凋亡)
- Neural communication and memory
- Relaxation of flat bronchial (支氣管) and gastrointestinal muscles (胃腸肌肉)
- Hormonal (激素) and sex functions
- Anti-microbial (抗微生物) and anti-tumor (抗腫瘤) defense
- Play an important role in tumor growth, immunodeficiency (免疫缺陷), cardiovascular (心血管), liver (肝), gastrointestinal tract (胃腸道) disease

NO treatment of wound pathologies





Before treatment



21st day of NO-therapy (10 seances)



After 2 months of NO-therapy

- Decrease in the trophic ulcer (營養性潰瘍) area:
 - Traditional treatment methods: 0.7% per day
 - NO treatment methods: 1.7% per day

G. Fridman, *et al.*, Plasma Process. Polym., **5**, 503 (2008) Plasma medicine, by Alexander Fridman and Gary Friedman

NO treatment of wound pathologies





Before treatment



After 4.5 months of NO-therapy (3 courses; 12 seances per course)

G. Fridman, *et al.*, Plasma Process. Polym., **5**, 503 (2008) Plasma medicine, by Alexander Fridman and Gary Friedman

Outline




- Example of several plasma discharges for plasma medicine
- Living tissue sterilization
- Blood coagulation
- Nitrogen oxide (NO) treatment
- Non-thermal plasma treatment of melanoma skin cancer
- Skin regeneration
- Egg sterilization
- Facemask regeneration

Non-thermal plasma treatment of melanoma skin cancer

(黑色素瘤皮膚癌)



- Melanoma cancer cell line (ATCC A2-58) was used
- ~1.5x10⁶ per dish








Figure 22. Results of FE-DBD treatment of melanoma cancer cells: Control, 5, 10, 20, and 30 s, counted 1, 3, and 24 h post-treatment. [27]

SKH-1 hairless mouse is treated with parallel plate electrode under isoflurane inhalation anesthesia








Melanoma shrinks after the treatment






- Day 0-3: 3 applications of 100 pulses (300 ns, 40 kv/cm, 0.5 Hz), 30 min apart
- Day 4: single application using 5 mm diameter parallel plate electrode

 Biochem Biophys Res Commun. 2006 May 5; 343(2): 351–360.

Electric field of 20 kV/cm is needed to treat Melanoma






Outline



- Example of several plasma discharges for plasma medicine
- Living tissue sterilization
- Blood coagulation
- Nitrogen oxide (NO) treatment
- Non-thermal plasma treatment of melanoma skin cancer
- Skin regeneration
- Egg sterilization
- Facemask regeneration

Plasma skin regeneration (PSR) is a novel skin treatment device





- PSR provides 1-2 J or 3-4 J per pulse for lower or higher power, respectively
- The skin is damaged slightly by the nitrogen plasma jet
- Skin regeneration is stimulated
- Local anesthetic (麻藥) is required and a systemic anesthetic, administered orally is recommended
- Ablative-like effect, similar to that of laser skin resurfacing can also be achieved, but with higher doses

Zones of the face and associated treatment energy settings







This particular patient-rated improvement in overall skin rejuvenation was 85%





 Patients reported minimal discomfort following the procedure and reported over 60% improvement in their skin condition

Outline



- Example of several plasma discharges for plasma medicine
- Living tissue sterilization
- Blood coagulation
- Nitrogen oxide (NO) treatment
- Non-thermal plasma treatment of melanoma skin cancer
- Skin regeneration
- Egg sterilization
- Facemask regeneration

Atmospheric-Pressure Plasma sterilization 99.9999% bacteria on surfaces of eggs





https://www.itri.org.tw/chi/Content/Publications/contents.aspx?Sit eID=1&MmmID=2000&MSid=745416417706673311


Outline



- Example of several plasma discharges for plasma medicine
- Living tissue sterilization
- Blood coagulation
- Nitrogen oxide (NO) treatment
- Non-thermal plasma treatment of melanoma skin cancer
- Skin regeneration
- Egg sterilization
- Facemask regeneration

A face mask do restrict the air flow from the mouth and the nose





Coughing over one breath w/o mask.


Coughing over a longer periods of time w/o mask.

Coughing over one breath w/ mask.

Talking w/o mask.

Wearing face mask can reduce the Covid-19 transmission probability significantly






Plasma can provide good surface treatment with low temperature



Treatment	Surface treatment level	Depth	Temperature	Cost
Chemical	Large	Deep	Room temperature ~200 °C	Medium
Heat	Only oxidizing	Deep	High temperature	Cheap
Radiation	Small	Whole sample	High temperature	Expensive
Plasma	Large	Surface	Room temperature ~100 °C	Cheap ~ Medium


 Atmospheric plasma can generate radicals, ozone, reactive oxygen/nitrogen/NH (ROS · RONS), UV light, electrons, charged particles.

Plasma can generate ROS and RONS




The temperature of the mask under plasma treamtment is below 40 ℃





The surface quality of the face mask was not influenced by the plasma treament






The growth of the bacteria on the face mask was suppressed





Disinfected (a) 24 hr

Without plasma treatment on used mask




With plasma treated on used mask





(e)




48 hr

DBD plasma demonstration







Show video.

DBD plasma can be generated between the finger and the dielectric layer



